

| Plant Name | Scientific | Common | Family |
|------------|------------|--------|--------|
|------------|------------|--------|--------|

|                              |  |                         |           |
|------------------------------|--|-------------------------|-----------|
| <i>Agastache</i> 'Pstessene' |  | CORONADO® Red<br>Hyssop | Lamiaceae |
|------------------------------|--|-------------------------|-----------|

| Propagation Method | Native to |
|--------------------|-----------|
|--------------------|-----------|

|          |                   |
|----------|-------------------|
| cuttings | Cultivated hybrid |
|----------|-------------------|

| Treatments (Suggested procedures) |
|-----------------------------------|
|-----------------------------------|

Cuttings should be taken from non-flowering shoots and should have at least 2 nodes, one above the media and one below. A talc based rooting hormone can improve rooting percentages. Stick cuttings in a well draining media under intermittent mist for the first 3-5 days.

| Collection Timing and Strategies |
|----------------------------------|
|----------------------------------|

Any time the plant is producing non-flowering stems.

| Cultural Information |
|----------------------|
|----------------------|

| Greenhouse conditions |
|-----------------------|
|-----------------------|

Start at 70F and 70% humidity, will typically callus in 2 weeks. Gradually, reduce humidity and temperature. Fertilizer levels should be medium. Normally, pinch plants between day 21 to 28.

| Garden conditions |
|-------------------|
|-------------------|

Plants are quite adaptable to many situations, but prefer full sun and moderate to dry, well draining soils.

| Other Expert Advice |
|---------------------|
|---------------------|

Florel at 500ppm will help reduce buds and blooms and promote more lateral growth.