

Plant Name	Scientific	Common	Family
------------	------------	--------	--------

<i>Delosperma</i> 'Psfave'		LAVENDER ICE ICE PLANT	Aizoaceae
----------------------------	--	------------------------	-----------

Propagation Method	Native to
--------------------	-----------

Cuttings	Cultivated hybrid
----------	-------------------

Treatments (Suggested procedures)

Treat 1½ - 2" tip cuttings with basal dip in 0.1% IBA rooting talc or 1:10 liquid rooting hormone. Stick cuttings in a well-drained medium and place under intermittent mist. Root initiation takes place in about 5 - 7days, and plugs are ready to transplant in 6-8 weeks.

Collection Timing and Strategies

Treat 1½ - 2" tip cuttings with basal dip in 0.1% IBA rooting talc or 1:10 liquid rooting hormone. Stick cuttings in a well-drained medium and place under intermittent mist. Root initiation takes place in about 5 - 7days, and plugs are ready to transplant in 6-8 weeks.

Cultural Information

Greenhouse conditions

Treat 1½ - 2" tip cuttings with basal dip in 0.1% IBA rooting talc or 1:10 liquid rooting hormone. Stick cuttings in a well-drained medium and place under intermittent mist. Root initiation takes place in about 5 - 7days, and plugs are ready to transplant in 6-8 weeks.

Garden conditions

Grows best in full sun to partial shade in clay, loam, or sandy soils. Requires moderate soil moisture, and is not as xeric as some succulents. Blooms from spring to fall.

Other Expert Advice

Well-drained soil is particularly important in the winter. Plants that are allowed to remain wet in winter are likely to rot.