

Plant Name	Scientific	Common	Family	Date Reviewed 03/24/20
<i>Delosperma</i> 'Psfave'		LAVENDER ICE ICE PLANT	Aizoaceae	
Propagation Method		Native to		
Cuttings		Cultivated hybrid		
Treatments (Suggested procedures)				
Treat 1½ - 2" tip cuttings with basal dip in 0.1% IBA rooting talc or 1:10 liquid rooting hormone. Stick cuttings in a well-drained medium and place under intermittent mist. Root initiation takes place in about 5 - 7days, and plugs are ready to transplant in 6-8 weeks.				
Collection Timing and Strategies				
Treat 1½ - 2" tip cuttings with basal dip in 0.1% IBA rooting talc or 1:10 liquid rooting hormone. Stick cuttings in a well-drained medium and place under intermittent mist. Root initiation takes place in about 5 - 7days, and plugs are ready to transplant in 6-8 weeks.				
Cultural Information				
Greenhouse conditions				
Treat 1½ - 2" tip cuttings with basal dip in 0.1% IBA rooting talc or 1:10 liquid rooting hormone. Stick cuttings in a well-drained medium and place under intermittent mist. Root initiation takes place in about 5 - 7days, and plugs are ready to transplant in 6-8 weeks.				
Garden conditions				
Grows best in full sun to partial shade in clay, loam, or sandy soils. Requires moderate soil moisture, and is not as xeric as some succulents. Blooms from spring to fall.				
Other Expert Advice				
Well-drained soil is particularly important in the winter. Plants that are allowed to remain wet in winter are likely to rot.				