

Plant Name	Scientific	Common	Family	Date Reviewed 03/24/20
Eriogonum umbellatum v. aureum 'Psdowns'		KANNAH CREEK® Buckwheat	Polygonaceae	
Propagation Method		Native to		
Seed or Cuttings		Western U.S.-particularly Colorado, Idaho, Utah, Nevada and Wyoming		
Treatments (Suggested procedures)				
Seeds- Seeds will germinate about 70% with no pre-treatment. Seeds require light to germinate do not cover seeds. Cuttings- Tip cuttings will root readily while plants are actively growing. Use a 1:10 liquid hormone treatment. Place cuttings under intermittent mist for 48 hours then hand water as needed. 3:1 perlite to peat media.				
Collection Timing and Strategies				
Seeds mature from the end of August through October. Cuttings can be taken May through September. Remove any flowering scapes on vegetative material.				
Cultural Information				
Greenhouse conditions				
Plants are susceptible to rot and should be kept on the dry side. Plants respond well to low amounts of liquid fertilizer.				
Garden conditions				
Plants are tolerant of a wide range of conditions, but perform best in full sun and in well-drained soil.				
Other Expert Advice				
Grows from low elevation plains deserts to sub alpine zones, up to 10,000 feet. Naturally grows in sand and gravel, rocky flats and slopes in mixed grassland and sagebrush areas. Also occurs in pinion, juniper and montane conifer woodlands.				